



Trochleoplasty and Tibial Tubercle Osteotomy (TTO) Post Operative Guidelines

INTRODUCTION

This is a general guide to rehabilitation for patients who have undergone a Trochleoplasty and TTO procedure by Mr John Jeffery, Consultant Orthopaedic Surgeon and Children's Knee Specialist.

Please note, when it comes to rehab, one size does not fit all. Rehab is physio led and may differ from these guidelines as a result of therapist preference, patient progress and local practices. Please check the post-operative note for any restrictions or variations from these guidelines.

GUIDANCE

The time scales are an approximate guide and may be altered depending on various factors such as pain, swelling and muscle control. The patient's management should be tailored to meet individual objectives.

PHASE 1: Extension (week 0-2)

Immediate post-operative stage.

Primary aims: Control pain and swelling, achieve full extension and quadriceps activation.

- Active and active-assisted knee flexion out of brace
- Static and inner range quadriceps exercises, Straight leg raise taught (test rather than exercise)
- Mobilise weight-bearing as tolerated with crutches in a brace locked in extension
- Ankle dorsiflexion/plantarflexion exercises, including weightbearing calf stretches
- Swelling management

Cautions:

	● Knee brace must be on and locked straight whenever mobilising - can be unlocked or removed when not mobilising
PHASE 2: Flexion (week 2-6) Osteotomy healing. Primary aims: Maintain full extension and achieve >90 degrees flexion. Wean off brace and crutches by 6 weeks.	● Mobilise weight-bearing as tolerated with brace unlocked and crutches ● Discard brace and crutches as soon as able ● Swelling management ● Scar management following wound review — soft tissue techniques to rectus femoris gentle stretching, concentric and eccentric exercises ● Start basic proprioception, balance and coordination training ● Consider core and hip stability exercises ● Open and closed chain knee flexion exercises ● Gait re-education; sit to stand, stair re-education — encourage incorporation into ADL ● Education regarding rehabilitation, and what to expect at each milestone. Address any fear-avoidance issues—reiterate the importance of the patient taking responsibility for increasing ROM and function Precautions: ● No resisted open-chain quads due to osteotomy site ● Avoid oversteering fixation/osteotomy with overpressure into flexion Contraindications
PHASE 3: Strength (week 6-12) Osteotomy consolidation. Primary aims: Achieve full range of motion. Introduce low-impact cardiovascular exercises. Build proximal strength and control.	● Exercises need to be tailored to patient's functional aim ● Once 100° flexion is achieved and has minimal swelling, can start using a stationary bike on minimal resistance ● Progressive closed chain, eccentric control exercises ● Road cycling allowed in flat pedals only ● Cardiovascular fitness ● Proprioceptive exercises — add controlled rotational exercises ● Swimming—freestyle and pool walking Precautions: ● If not progressing flexion beyond 90 degrees then focus on rectus femoris stretches ● No resisted open-chain quads or impact activity until osteotomy united ● No breaststroke until 3 months

PHASE 4: Sport-specific rehab (>12 weeks)

By this stage the osteotomy has consolidated.

Primary aims: Introduce high-impact cardiovascular exercises. Increase fitness and work towards normal gait in sprinting and good control of cutting, pivoting, stopping and starting.

- Increase fitness
- Introduction of impact work — only if an X-ray has confirmed osteotomy union and the patient has a full range of motion with eccentric quadriceps control and correct alignment.
- Gradual increase in resisted open-chain/closed chain quadriceps (avoid pain)
- Continue with proprioceptive training—increase rotational control
- Initiate running—gradual paced change of terrain/gradient and duration
- Progressive introduction of dynamic activity:
 - jumping/hopping (start on the trampette, emphasis on alignment of both push off and land)
 - change of direction; start single direction and progress to cutting, multidirectional and pivoting
 - stopping/starting and acceleration/deceleration
 - backwards running

RETURN TO SPORT

Final decision on return to sport is up to the individual physiotherapist after assessment of the patient's progress and milestones. Return to training and non-match play can be expected at about 4 months post op.

We do not recommend returning to competitive/contact pivoting sport (i.e. match play) until 6 months post-surgery and this should be built up to in a graded fashion as with any rehabilitation. We do not have a preferred return-to-play criteria to use but example criteria include:

- >80% hop height, length and cross over
- >80% strength of non-involved limb
- Confidence in knee
- Awareness of safe positioning of limb and cutting/landing technique (see poweruptoplay.org)

FUNCTIONAL MILESTONES SUMMARY

Activity	Time Scale
High-impact exercise (jogging)	3 months
Sport-specific drills	4 months
Non-match sport	5 months
Competitive sports	6 months

RED FLAGS

The following should prompt urgent referral back to clinic:

- Signs of infection
- Thrombosis
- Persistent stiffness >8 weeks

Mr John Jeffery

MBBS BSc FRCS (Tr & Orth) PGDip (Med Ed)



CHILDREN'S KNEE SPECIALIST

Consultant Paediatric Orthopaedic Surgeon

Secretary: Tara James

Tel: 0117 9804052

Email: tara.james@spirehealthcare.com